

REVIEW

Heart rate variability in male rats

Pavol Švorc Jr¹ | Soňa Grešová²  | Pavol Švorc^{1,2}

¹Department of Physiology and Pathophysiology, Faculty of Medicine, Ostrava University, Ostrava, Czech Republic

²Department of Physiology, Faculty of Medicine, University PJ Safarika, Tr. SNP 1, Kosice 04001, Slovak Republic

Correspondence

Soňa Grešová, Department of Physiology, Faculty of Medicine, University PJ Safarika, Tr. SNP 1, Kosice 04001, Slovak Republic.
Email: sona.gresova@upjs.sk

Abstract

The cardiovascular system is primarily controlled by the autonomic nervous system, and any changes in sympathetic or parasympathetic activity also have an impact on myocardial activity. Heart rate variability (HRV) is a readily available metric used to assess heart rate control by the autonomic nervous system. HRV can provide information about neural (parasympathetic, sympathetic, reflex) and humoral (hormones, thermoregulation) control of myocardial activity. Because there are no relevant reference values for HRV parameters in rats in the scientific literature, all experimental results are only interpreted on the basis of changes from currently measured control or baseline HRV values, which are, however, significantly different in individual studies. Considering the significant variability of published HRV data, the present study focused primarily on comparing control or baseline HRV values under different conditions in in vivo experiments involving rats. The aim of the study was therefore to assess whether there are differences in the starting values before the experiment itself.

KEYWORDS

heart rate variability, in vivo experiments, rats

1 | INTRODUCTION

There is an existing problem in determining “normal” values of heart rate variability (HRV) parameters in frequency-domain or time-domain analyses. Relatively few investigators have used HRV parameters only in terms of changes after a given intervention, derived from baseline values, while not accounting for the fact that the evaluated parameters—or their extent—can be altered already at the beginning of the experiment. Such an approach is understandable given that no standard reference values for HRV in rats have been reported. Chizh (2015) also draws attention to this problem, where he points out the difference in the frequency ranges of various authors, from which spectral components were evaluated. This reflects the

considerable variability in baseline or control values at the evaluation of the frequency-domain or time-domain HRV parameters themselves. In the present article, we focus on showing differences in control or baseline values of HRV parameters under different conditions in experiments involving rats, which can influence the accuracy of the interpretation of the results and the conclusions drawn.

2 | HRV IN EXPERIMENTS INVOLVING RATS

The concept that HRV in rodents similarly reflects cardiovascular regulation in humans is supported. Low or missing resting vagus nerve tone in these species, however,

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Physiological Reports* published by Wiley Periodicals LLC on behalf of The Physiological Society and the American Physiological Society.

can alter the quality and degree of HRV response after a given intervention. Although the origin of variation in the frequency-domain (Kuwahara et al., 1994) and time-domain parameters of HRV (Aubert et al., 1999) appears to be analogous to humans, the degree of influence of the parasympathetic division may be greater in rats, as supported by several previous studies (Cerutti et al., 1991; Japundzic et al., 1990).

On the contrary, (Rowan III et al., 2007). queried to what extent changes in HRV can be used as a marker of increased or decreased sensitivity to a given intervention in rodent studies. Perhaps the most appropriate interpretation of most short-term studies involving rodents is simply that intervention(s) may produce a systemic effect that is mediated either by the nervous system or by the heart or blood vessels, but this is pure speculation and not necessarily supported by concrete evidence. Nevertheless, HRV may still play an important mechanistic role in such research. To better understand the prognostic utility of HRV together with other electrocardiographic (ECG) endpoints, it is necessary to proceed from carefully controlled experiments with known outcomes (Carnevali et al., 2019; Krüger et al., 2000; Sanyal et al., 2002). Therefore, the development of new analytical techniques that connect temporal trends in cardiovascular parameters with pathological conditions is necessary—sometimes even essential—and may provide a basis for future studies using rats with large individual differences in vagus-mediated HRV.

3 | HRV: EVALUATION

In the study, we included the values of frequency and time parameters of HRV from studies retrieved from a literature search of the Web of Science database using the keywords “HRV in rat.” HRV parameters that were reported as baseline or control values were summed and means were calculated. In some studies, HRV parameters were reported from the time domain and in others from the frequency domain and, in some cases, HRV was assessed using both methods. Most of the studies described changes in HRV only in graphical form without reporting actual numerical values, or only changes in HRV were noted.

3.1 | Evaluated groups

Female, juvenile, and old animals were excluded from the study. The age of the rats ranged from 3 to 4 months (i.e., sexually mature male rats). Rat strain and telemetry equipment for HRV measurements were not taken into account. The length of the records from which HRV was

evaluated was only described in some methodologies; as such, taking this factor into account was highly problematic. For a clearer comparison of HRV parameters, we created the following groups: *control group*—awake or freely moving animals, HRV was measured telemetrically before the experiment itself; and *sham group*—awake or freely moving animals, in which the animals, after the introduction of telemetric sensing, underwent surgical preparation for the ongoing experiment, but without subsequent targeted experimental intervention.

The most frequently used frequency-domain parameters (LF power, HF power, LF/HF ratio, LFnu, HFnu and total spectral power (TSP)) and time-domain parameters (RR interval, SDNN and rMSSD) from selected studies were evaluated and compared.

3.2 | Statistical analysis

Data are expressed as mean \pm standard deviation (SD). Data were analyzed using InStat (GraphPad). The Tukey–Kramer test was used to compare data from the groups, and differences with $p < 0.05$ were considered to be statistically significant. Correlations were calculated as correlation coefficients, in which the coefficient was statistically significant in the range of -0.4 to -1 and from $+0.4$ to $+1$.

4 | HRV: CONTROL (BASELINE) VALUES FROM FREQUENCY-DOMAIN ANALYSIS

Experimental designs are always based on control or reference values, which fully apply to the activities of the autonomic nervous system. The autonomic nervous system of the rat is very sensitive to various external stimuli and/or those from the internal environment, which can significantly alter the activity of the autonomic nervous system at the beginning of an experiment. Therefore, it is necessary to carefully consider preparation of the animal for the experiment itself and to determine which values to define as the starting point (i.e., baseline) if the desire is to assess not only changes in the autonomic nervous system themselves but also changes in relation to the monitored parameter.

In in vivo experiments involving awake or freely moving animals, baseline values are used from sham groups, when the animals, in addition to the implantation of electrodes for continuous ECG recording, are also subjected to other surgical procedures as preparation for the further course of the experiment, but without subsequent targeted experimental intervention. For example, preparation for coronary or femoral artery occlusion (Aires et al., 2017; De La Fuente et al., 2013; Nobre

et al., 2006), catheterization (Beckers et al., 2006; Blanco et al., 2015; Fiorino et al., 2012; Krüger et al., 1997, 2000; Mostarda et al., 2009; Müller-Ribeiro et al., 2017; Quagliotto et al., 2015; Ramaekers et al., 2002; Ribeiro et al., 2021; Sallam et al., 2016, 2017; Shi et al., 2020; Simoes et al., 2016), ligation (Lima et al., 2018; Maida et al., 2017; Ruiz et al., 2014; Wang et al., 2016), bile duct ligation (Haddadian et al., 2013), cannulation (Beltrán et al., 2020; Da Silva et al., 2002; Neto et al., 2017; Sant'Ana et al., 2011), implantation of stimulators (Dai et al., 2020; Domingos-Souza et al., 2021), or if they have already been administered substances beforehand (Lo Giudice et al., 2002).

Although the animals used in these experiments undergo a recovery phase (when antibiotics and analgesics are administered), it is not reported anywhere whether the autonomic nervous system modulation over the heart “returns to normal” after recovery from surgical interventions or whether the changes persist from a longer-term perspective. The overall recovery time after surgery can be a problem. In the methodologies, this time varies from 1 hour to 3–4 weeks and is likely dependent on the severity of the intervention.

In control group, HRV was measured telemetrically before the experiment itself in awake and freely moving male animals (Carll et al., 2012; Choudhary et al., 2018; Couderc et al., 2002; Fazan Jr. et al., 2015; Hazari et al., 2021; Imai et al., 2008; Koizumi et al., 2011; Lamb et al., 2012; Lin et al., 2016; Mamalyga, 2013; Mangin et al., 1998; Pereira-Junior et al., 2006, 2010; Saalfeld & Spear, 2014; Shi et al., 2017; Soler et al., 2018; Towa et al., 2004; Tsai et al., 2020; Yang et al., 2019; Zajackowski et al., 2014). If we assume that the location of implantation (subcutaneous on the back or intra-abdominal) and the length of recovery do not play a role, then theoretically we can assume that these values could approach the “missing” reference values, although unfortunately, only for males and at the given time of measurement. On the contrary, also during electrode implantation itself, the animals are anesthetized using different types of anesthetics (pentobarbital, ketamine/xylazine, urethane, and isoflurane), while general anesthesia is known to suppress HRV in human (Matchett & Wood, 2014) and also in animals (Mäenpää et al., 2007).

To determine whether there are any differences between sham and control groups, we focused only on some evaluated parameters from the frequency-domain analysis. Differences in baseline values are reported in Table 1 and Figure 1. In the control group, although the HF power (representing the modulation of the parasympathetic) prevails, the LF/HF ratio points to a state in which the sympathetic is significantly dominant. The dominant sympathetic effect is also indicated by the value of LFnu (normalized unit represents the relative

TABLE 1 Mean \pm SD values of frequency-domain HRV parameters from conscious male rats.

Group	VLF power (ms^2)	LF power (ms^2)	HF power (ms^2)	LF/HF	LFnu	HFnu	TSP (ms^2)
Sham	12.4 \pm 12.6 (n = 6)	3.05 \pm 3.06 (n = 34)	8.04 \pm 9.03 (n = 36)	0.75 \pm 0.8 (n = 33)	23.6 \pm 18.8 (n = 26)	57.3 \pm 23.5 (n = 24)	20.98 \pm 8.4 (n = 12)
Control	16.46 \pm 13.34 (n = 8)	3.22 \pm 2.36 (n = 18)	5.45 \pm 3.49 (n = 19)	1.49 \pm 1.46 (n = 25) $p < 0.02$	50.9 \pm 30.6 (n = 4)	48.7 \pm 30.7 (n = 4)	24.3 \pm 17.9 (n = 8)

Note: Sham—animals which, in addition to the implantation of electrodes for continuous electrocardiographic recording, also underwent other surgical interventions but without targeted experimental intervention; Control—animals that only underwent implantation of electrodes; VLF—is not as well characterized but is believed to reflect circadian inputs to the heart; LF—sympathetic and parasympathetic activities or baroreceptor activity; HF—parasympathetic activity; LF/HF ratio—reflects sympathovagal balance; LFnu, HFnu—normalized units; TSP—presents total HRV; n = number of measurements from which the average value was calculated. $p < 0.02$ statistically significant difference between sham and control groups.

Abbreviations: HF, high frequency; LF, low frequency; TSP, total spectral power; VLF, very low frequency.




FIGURE 1 Mean \pm SD values of individual frequency HRV parameters from conscious male rats (sham and control). VLF, very low frequency of HRV (VLF-sham $n=6$; VLF-control $n=8$); LF, low frequency of HRV (LF-sham $n=34$; LF-control $n=18$); HF, high frequency of HRV (HF-sham $n=36$; HF-control $n=19$); TSP, total spectral power of HRV (TSP-sham $n=12$; TSP-control $n=8$).




FIGURE 2 Mean \pm SD values of LFnu and HFnu conscious male rats (sham vs. control). LFnu (LFnu-sham $n=26$; LFnu-control $n=4$) and HFnu (HFnu-sham $n=24$; HFnu-control $n=4$)—normalized units represent the relative value of the HF and LF components of HRV in relation to TSP (after subtracting VLF power).

value of the LF component of HRV in relation to TSP), which is higher than HFnu (normalized unit represents the relative value of the HF component of HRV in relation to TSP), in contrast to the sham group, in which LFnu is lower than HFnu (Table 1, Figure 2). Based on averaged data, it appears that total HRV (presented by TSP) is reduced in the sham group, although not significantly. Could this mean that the autonomic nervous system in a rat sham group is less sensitive to various manipulations or interventions than the control group?

However, this only occurs during the light (i.e., inactive) period of the rat regimen day, when the experiments were performed.

4.1 | HRV: control (baseline) values from time-domain analysis

Similar to investigations that analyzed changes in autonomic nervous system activity using frequency-domain analysis,

other studies have used time-domain analysis of HRV. As in frequency-domain studies, baseline values not only in sham groups were used (Abulaiti et al., 2011; De La Fuente et al., 2013; Houshmand et al., 2017; Krüger et al., 1997, 2000; Lima et al., 2018; Müller-Ribeiro et al., 2017; Sallam et al., 2016, 2017; Silva, Geraldini, et al., 2017; Silva, Silva, et al., 2017; Soler et al., 2018; Wang et al., 2016; Xu et al., 2020), but also from groups in which the values were recorded only before the experimental intervention itself and the animals did not undergo surgical preparation (control) (Barbier et al., 2006; Beckers et al., 2006; Beltrán et al., 2020; Couderc et al., 2002; Farraj et al., 2009; Fazan Jr. et al., 2015; Hashimoto et al., 1999; Hazari et al., 2021; Koresh et al., 2016; Lamb et al., 2012; Lin et al., 2016; Maida et al., 2017; Mangin et al., 1998; Mostarda et al., 2009; Pereira-Junior et al., 2010; Powell et al., 2021; Ramaekers et al., 2002; Ribeiro et al., 2021; Saalfeld & Spear, 2014; Scridon et al., 2012; Shi et al., 2017; Simoes et al., 2016; Zajączkowski et al., 2018).

TABLE 2 Mean (\pm SD) values of time-domain HRV parameters from conscious male rats.

Group	Mean RR (ms)	SDNN (ms)	rMSSD (ms)
Sham	157.6 \pm 14.5 (<i>n</i> = 15)	7.93 \pm 3.92 (<i>n</i> = 11)	4.89 \pm 1.93 (<i>n</i> = 7)
Control	170.7 \pm 23.2 (<i>n</i> = 13) <i>p</i> < 0.001	9.48 \pm 6.75 (<i>n</i> = 25)	4.99 \pm 2.24 (<i>n</i> = 23)

Note: Sham—animals which, in addition to the implantation of electrodes for continuous electrocardiographic recording, also underwent other surgical interventions, but without targeted experimental intervention; Control—animals that only underwent implantation of electrodes; *n* = number of measurements from which the average value was calculated. *p* < 0.001 statistically significant difference between sham and control groups.


From data obtained from individual studies involving male rats, significant differences (*p* < 0.001) have been found in the duration of RR intervals between the sham and control groups, while the duration of RR intervals varied depending on parasympathetic activity (rMSSD) in both groups (sham group, *r* = −0.58; control group, *r* = −0.96; Table 2). Based on comparison of the calculated correlation coefficients from both groups, we can only speculate that, in the control group, heart rate is practically and completely dependent on changes in parasympathetic activity, while in the sham group, changes in heart rate can also be the result of the action of other mechanisms. The contribution of the sympathetic nervous system to the duration of the RR intervals from the given data is unclear. Regarding the contribution of total HRV (SDNN) to the duration of RR intervals, a moderate significant dependence was found only in the control group (sham group, *r* = −0.15; control group, *r* = 0.49; Table 2, Figure 3).

5 | DISCUSSION AND CONCLUSION

Results from each of the cited studies are, of course, excellent. It is obvious in experimental studies that when evaluation of any changes in monitored parameters, it should be based on the control or baseline values. However, the question remains as to why there is so much variability in these values. A follow-up question would be to what extent the control or baseline values can be—or they are—changed already at the beginning of the experiment itself.

One of the possible problems of great variance HRV may be the nonuniformity in frequency ranges, from which VLF, LF, HF, and TSP are calculated (Table 3). There

FIGURE 3 Averaged values of SDNN and rMSSD from conscious male rats (sham and control). SDNN (ms) (SDNN-sham *n* = 11; SDNN-control *n* = 25)—a frequently used index of total HRV identical to TSP. rMSSD (ms) (rMSSD-sham *n* = 7; rMSSD-control *n* = 23)—higher rMSSD values indicate greater parasympathetic activity.



Author	VLF (Hz)	LF (Hz)	HF (Hz)	TSP (Hz)
Krüger et al. (2000, 1997)		0.5–0.8	≥0.8	
Mangin et al. (1998)		0.2–1	1–3	
Zajaczkowski et al. (2014, 2018)		0.2–0.8	0.8–2.5	
Pereira-Junior et al. (2006, 2010)				
Fazan Jr. et al. (2015)	<0.2	0.2–0.8	0.8–2	
Shi et al. (2017)		0.04–0.6	0.6–2.4	
Mamalyga (2013)	0.02–0.8			
Carll et al. (2012)		0.2–0.8	0.8–2	
Koizumi et al. (2011)		0.3–0.8	0.9–3.3	
Imai et al. (2008)		0.04–1	1–3	
Towa et al. (2004)				
Shi et al. (2020)	0.01–0.04	0.1–1	1–3	
Ramaekers et al. (2002)		0.2–0.7	0.8–2.5	
Beckers et al. (2006)				
Couderc et al. (2002)		0.1–0.7	0.7–2	
Beltrán et al. (2020)		0.04–0.6	0.6–2.4	
De La Fuente et al. (2013)	0–0.2	0.2–0.8	0.8–4	
Mostarda et al. (2009)	0–0.2	0.2–0.8	0.8–3	
Quagliotto et al. (2015)				
Blanco et al. (2015)				
Simoes et al. (2016)				
Müller-Ribeiro et al. (2017)				
Lo Giudice et al. (2002)	0.03–0.2	0.2–0.6	0.60–3	
Ribeiro et al. (2021)		0.2–0.8	0.8–3	
Aires et al. (2017)				
Dai et al. (2020)		0.3–0.8	0.8–4	
Wang et al. (2016)		0.1–0.6	0.6–3	
Sallam et al. (2016, 2017)		0.3–0.8	0.8–3	
Abulaiti et al. (2011)	0.003–0.04	0.04–0.2	0.2–0.4	<0.4
Lin et al. (2016)				
Lima et al. (2018)		0.04–0.2	0.2–0.4	
Chang et al. (2011)				
Yang et al. (2019)	0.02–0.2	0.2–0.6	0.6–3	0–3
Tsai et al. (2020)				
Hazari et al. (2021)		0.2–0.8	0.8–3.5	
Da Silva et al. (2002)		0.2–0.8	0.8–2.5	
Fiorino et al. (2012)		0.2–0.6	0.6–3	
Neto et al. (2017)	0.01–0.2	0.2–0.8	0.8–3	
Maida et al. (2017)				
Sant'Ana et al. (2011)	0.01–0.2	0.2–0.85	0.8–2.5	
Zhu et al. (2018)	0.003–0.04	0.04–0.2	0.2–0.4	0–0.5

TABLE 3 Values of frequency range from individual authors.

is also inconsistency in the use of units for HRV frequency parameters, because some investigators have described them in ms^2 , some in ms^2/Hz , some only in ms , and some not at all. It is assumed that algorithms for the calculation of single frequency and time parameters of HRV should be constant; as such, calculation from different frequency ranges can be one of the causes of variability, because

there are enormous differences in the reporting of values, which, in some studies, can differ by orders of magnitude (Chang et al., 2011; Fabiyi-Edebor, 2020; Malliani et al., 1991; Zhu et al., 2018).

Another problem may be the fact that many methodologies do not describe the interval duration from which individual HRV parameters were evaluated. Different

methods of HRV acquisition and evaluation, as well as sex, use of anesthesia during telemetry device implantation, recovery period, and time when the measurement was performed may also be problems (Švorc Jr. et al., 2023).

Analysis of baseline or control values reported in the cited studies clearly reflect wide disparity, and there are no “normal” or standardized reference values for HRV parameters in rats from frequency-domain or time-domain analyses of HRV obtained in a specifically defined time of measurement. Therefore, we believe that unequivocal conclusions should not be drawn, especially those regarding the autonomic nervous system and should, therefore, be offered cautiously.

AUTHOR CONTRIBUTIONS

Pavol Švorc proposed the original draft of the study. Pavol Švorc Jr and Soňa Grešová collected all data and performed data analysis. Pavol Švorc designed the first manuscript. Pavol Švorc and Soňa Grešová drafted the final manuscript. All authors made critical revisions and approved the final version of the manuscript. All authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

ACKNOWLEDGMENTS

The present study was supported by VEGA grant 1/0008/20 of the Ministry of Education of the Slovak Republic.

FUNDING INFORMATION

None.

CONFLICT OF INTEREST STATEMENT

No conflicts of interest, financial or otherwise, are declared by the authors.

DATA AVAILABILITY STATEMENT

Data were obtained from several scientific articles.

ORCID

Soňa Grešová  <https://orcid.org/0000-0001-5964-8184>

REFERENCES

- Abulaiti, A., Hu, D., Zhu, D., & Zhang, R. (2011). Influence of fastigial nucleus stimulation on heart rate variability of surgically induced myocardial infarction rats: Fastigial nucleus stimulation and autonomous nerve activity. *Heart and Vessels*, 26, 654–662.
- Aires, R., Pimentel, E. B., Forechi, L., Dantas, E. M., & Mill, J. G. (2017). Time course of changes in heart rate and blood pressure variability in rats with myocardial infarction. *Brazilian Journal of Medical and Biological Research*, 50, e5511.
- Aubert, A. E., Ramaekers, D., Beckers, F., Breem, R., Denef, C., van de Werf, F., & Ector, H. (1999). The analysis of heart rate variability in unrestrained rats. Validation of method and results. *Computer Methods and Programs in Biomedicine*, 60, 197–213.
- Barbier, J., Reland, S., Ville, N., Rannou-Bekono, F., Wong, S., & Carré, F. (2006). The effects of exercise training on myocardial adrenergic and muscarinic receptor. *Clinical Autonomic Research*, 16, 61–65.
- Beckers, F., Verheyden, B., Ramaekers, D., Swynghedauw, B., & Aubert, A. E. (2006). Effects of autonomic blockade on non-linear cardiovascular variability indices in rats. *Clinical and Experimental Pharmacology and Physiology*, 33, 431–439.
- Beltrán, A. R., Arce-Álvarez, A., Ramírez-Campillo, R., Vázquez-Muñoz, M., von Igel, M., Ramírez, M. A., Del Rio, R., & Andrade, D. C. (2020). Baroreflex modulation during acute high-altitude exposure in rats. *Frontiers in Physiology*, 11, 1049.
- Blanco, J. H. D., Gastaldi, A. C., Gardim, C. B., Araujo, J. E., Simoes, M. V., Oliviera, L. F. L., Carvalho, E. E. V., & Souza, H. C. D. (2015). Chronic cholinergic stimulation promotes changes in cardiovascular autonomic control in spontaneously hypertensive rats. *Autonomic Neuroscience*, 193, 97–103.
- Carll, A. P., Hazari, M. S., Perez, C. M., Krantz, Q. T., King, C. J., Winsett, D. W., Costa, D. L., & Farraj, A. K. (2012). Whole and particle-free diesel exhausts differentially affect cardiac electrophysiology, blood pressure, and autonomic balance in heart failure-prone rats. *Toxicological Sciences*, 128, 490–499.
- Carnevali, L., Statello, R., & Sgoifo, A. (2019). Resting heart rate variability predicts vulnerability to pharmacologically-induced ventricular arrhythmias in male rats. *Journal of Clinical Medicine*, 8, 655.
- Cerutti, C., Gustin, M. P., Paultre, C. Z., Lo, M., Julien, C., Vincent, M., & Sassard, J. (1991). Autonomic nervous system and cardiovascular variability in rats: A spectral analysis approach. *The American Journal of Physiology*, 261, H1292–H1299.
- Chang, Y. T., Wann, S. R., Wu, P. L., Hsieh, K. H., Lin, C. C., Huang, M. S., & Chang, H. T. (2011). Influence of age on heart rate variability during therapeutic hypothermia in a rat model. *Resuscitation*, 82, 1350–1354.
- Chizh, N. A. (2015). Parameters of spectral analysis of heart rate variability in rats. *Problems of Cryobiology and Cryomedicine*, 25(3), 235–245.
- Choudhary, A. K., Sundareswaran, L., & Devi, R. S. (2018). Effects of aspartame on the evaluation of electrophysiological responses in Wistar albino rats. *Journal of Taibah University for Science*, 10, 505–512.
- Couderc, J. P., Elder, A., Cox, C., Zareba, W., & Oberdorster, G. (2002). Limitations of power-spectrum and time-domain analysis of heart rate variability in short-term ECG recorded using telemetry in unrestrained rats. *Computers in Cardiology*, 29, 589–592.
- Da Silva, V. J. D., Viana, P. C. C., Alves, R. M., Fazan, R., Jr., Ruscone, T. G., Porta, A., Malliani, A., Salgado, H. C., & Montano, N. (2002). Intravenous amiodarone modifies autonomic balance and increases baroreflex sensitivity in conscious rats. *Autonomic Neuroscience*, 95, 88–96.

- Dai, F., Yin, J., & Chen, J. D. D. Z. (2020). Effects and mechanisms of vagal nerve stimulation on body weight in diet-induced obese rat. *Obesity Surgery*, 30, 948–956.
- De La Fuente, R. N., Rodrigues, B., Moraes-Silva, I. C., Souza, L. E., Sirvente, R., Mostarda, C., De Angelis, K., Soares, P. P., Lacchini, S., Consolim-Colombo, F., & Irigoyen, M. C. (2013). Cholinergic stimulation with pyridostigmine improves autonomic function in infarcted rats. *Clinical and Experimental Pharmacology & Physiology*, 40, 610–616.
- Domingos-Souza, G., Santos-Almeida, F. M., Meschiari, C. A., Ferreira, N. S., Pereira, C. A., Pestana-Oliviera, N., Prates-Costa, T. C., Tostes, R. C., White, C., & Fazan, R., Jr. (2021). The ability of baroreflex activation to improve blood pressure and resistance vessel function in spontaneously hypertensive rats is dependent on stimulation parameters. *Hypertension Research*, 44, 932–940.
- Fabiyyi-Edebor, T. D. (2020). Vitamin C ameliorated cardiac autonomic neuropathy in type 2 diabetic rats. *World Journal of Diabetes*, 11, 52–65.
- Farraj, A. K., Haykal-Coates, N., Winsett, D. W., Hazari, M. S., Carll, A. P., Rowan, W. H., III, Ledbetter, A. D., Cascio, W. E., & Costa, D. L. (2009). Increased nonconducted P-wave arrhythmias after a single oil fly ash inhalation exposure in hypertensive rats. *Environmental Health Perspectives*, 117, 709–715.
- Fazan, R., Jr., Silva, C. A. A., Oliveira, J. A. C., Salgado, H. C., Montano, N., & Garcia-Cairasco, N. (2015). Evaluation of cardiovascular risk factors in the Wistar audiogenic rat (WAR) strain. *PLoS ONE*, 10, e0129574.
- Fiorino, P., Evangelista, F. S., Santos, F., Magri, F. M. M., DeLorenzi, L. C. M. O. B., Ginoza, M., & Farah, V. (2012). The effect of green tea consumption on cardiometabolic alterations induced by experimental diabetes. *Experimental Diabetes Research*, 2012, 309231.
- Haddadian, Z., Eftekhari, G., Mazloom, R., Jazaeri, F., Dehpour, A. R., & Mani, A. R. (2013). Effect of endotoxin on heart rate dynamics in rats with cirrhosis. *Autonomic Neuroscience*, 177, 104–113.
- Hashimoto, M., Kuwahara, M., Tsubone, H., & Sugano, S. (1999). Diurnal variation of autonomic nervous activity in the rat—investigation by power spectral analysis of heart rate variability. *Journal of Electrocardiology*, 32, 167–171.
- Hazari, M. S., Phillips, K., Stratford, K. M., Khan, M., Thompson, L., Oshiro, W., Hudson, G., Herr, D. W., & Farraj, A. K. (2021). Exposure to intermittent noise exacerbates the cardiovascular response of Wistar-Kyoto rats to ozone inhalation and arrhythmogenic challenge. *Cardiovascular Toxicology*, 21, 336–348.
- Houshmand, F., Faghihi, M., Imani, A., & Kheiri, S. (2017). Effect of different doses of oxytocin on cardiac electrophysiology and arrhythmias induced by ischemia. *Journal of Advanced Pharmaceutical Technology and Research*, 8, 131–137.
- Imai, K., Ariga, H., Chen, C., Mantyh, C., Pappas, T. N., & Takahashi, T. (2008). Effects of electroacupuncture on gastric motility and heart rate variability in conscious rats. *Autonomic Neuroscience: Basic and Clinical*, 138, 91–98.
- Japundzic, N., Grichois, M. L., Zitoun, P., Laude, D., & Elghozi, J. L. (1990). Spectral analysis of blood pressure and heart rate in conscious rats: Effects of autonomic blockers. *Journal of the Autonomic Nervous System*, 30, 91–100.
- Koizumi, S., Minamisawa, S., Sasaguri, K., Onozuka, M., Sato, S., & Ono, Y. (2011). Chewing reduces sympathetic nervous response to stress and prevents poststress arrhythmias in rats. *American Journal of Physiology. Heart and Circulatory Physiology*, 301, H1551–H1558.
- Koreish, O., Kaplan, Z., Zohar, J., Matar, M. A., Geva, A. B., & Cohen, H. (2016). Distinctive cardiac autonomic dysfunction following stress exposure in both sexes in an animal model of PTSD. *Behavioural Brain Research*, 308, 128–142.
- Krüger, C., Kalenka, A., Haunstetter, A., Schweizer, M., Maier, C., Rühle, U., Ehmke, H., Kübler, W., & Haass, M. (1997). Baroreflex sensitivity and heart rate variability in conscious rats with myocardial infarction. *The American Journal of Physiology*, 273, H2240–H2247.
- Krüger, C., Landerer, V., Zugck, C., Ehmke, H., Kübler, W., & Haass, M. (2000). The bradycardic agent zatebradine enhances baroreflex sensitivity and heart rate variability in rats early after myocardial infarction. *Cardiovascular Research*, 45, 900–912.
- Kuwahara, M., Yayou, K., Ishii, K., Hashimoto, S., Tsubone, H., & Sugano, S. (1994). Power spectral analysis of heart rate variability as a new method for assessing autonomic activity in the rat. *Journal of Electrocardiology*, 27, 333–337.
- Lamb, C. M., Hazari, M. S., Haykal-Coates, N., Carll, A. P., Krantz, Q. T., King, C., Winsett, D. W., Cascio, W. E., Costa, D. L., & Farraj, A. K. (2012). Divergent electrocardiographic responses to whole and particle-free diesel exhaust inhalation in spontaneously hypertensive rats. *Toxicological Sciences*, 125, 558–568.
- Lima, S. E. B., de Oliveira, L. C. S., Cardoso, G. S., Telles, P. V. N., Lima, L. C., E Sousa, J. F. R., Araújo, R. P. N., Oliveira, A. P., Santos, R. F., Santos, A. A., & Silva, M. T. B. (2018). Moderate-intensity exercise and renin angiotensin system blockade improve the renovascular hypertension (2K1C)-induced gastric dysmotility in rats. *Life Sciences*, 210, 55–64.
- Lin, T. T., Sung, Y. L., Wu, C. E., Zhang, H., Liu, Y. B., & Lin, S. F. (2016). Proarrhythmic risk and determinants of cardiac autonomic dysfunction in collagen-induced arthritis rats. *BMC Musculoskeletal Disorders*, 17, 491.
- Lo Giudice, P., Careddu, A., Magni, G., Quagliata, T., Pacifici, L., & Carminati, P. (2002). Autonomic neuropathy in streptozotocin diabetic rats: Effect of acetyl-L-carnitine. *Diabetes Research and Clinical Practice*, 56, 173–180.
- Mäenpää, M., Penttilä, J., Laitio, T., Kaisti, K., Kuusela, T., Hinkka, S., & Scheinin, H. (2007). The effects of surgical levels of sevoflurane and propofol anaesthesia on heart rate variability. *European Journal of Anaesthesiology*, 24, 626–633.
- Maida, K. D., Gastaldi, A. C., Faccioli, T. P., Araújo, J. E., & de Souza, H. C. D. (2017). Physical training associated with enalapril but not to losartan, results in better cardiovascular autonomic effects. *Autonomic Neuroscience: Basic and Clinical*, 203, 33–40.
- Malliani, A., Pagani, M., Lombardi, F., & Cerutti, S. (1991). Cardiovascular neural regulation explored in the frequency domain. *Circulation*, 84, 482–492.
- Mamalyga, M. L. (2013). Heart rate regulation at different levels of convulsive readiness. *Bulletin of Experimental Biology and Medicine*, 155, 425–428.
- Mangin, L., Swynghedauw, B., Benis, A., Thibault, N., Lerebours, G., & Carré, F. (1998). Relationships between heart rate and heart rate variability: Study in conscious rats. *Journal of Cardiovascular Pharmacology*, 32, 601–607.
- Matchett, G., & Wood, P. (2014). General anesthesia suppresses normal heart rate variability in humans. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 24(2), 23129.

- Mostarda, C., Rogow, A., Silva, I. C. M., De La Fuente, R. N., Jorge, L., Rodrigues, B., Heeren, M. V., Caldini, E. G., De Angelis, K., & CláudiaIrigoyen, M. (2009). Benefits of exercise training in diabetic rats persist after three weeks of detraining. *Autonomic Neuroscience*, 145, 11–16.
- Müller-Ribeiro, F. C., Wanner, S. P., Santos, W. H. M., Malheiros-Lima, M. R., Fonseca, I. A. T., Coimbra, C. C., & Pires, W. (2017). Changes in systolic arterial pressure variability are associated with the decreased aerobic performance of rats subjected to physical exercise in the heat. *Journal of Thermal Biology*, 63, 31–40.
- Neto, O. B., Sordi, C. C., Mota, G. R., Marocolo, M., Chrighuer, R. S., & Silva, V. J. D. (2017). Exercise training improves hypertension-induced autonomic dysfunction without influencing properties of peripheral cardiac vagus nerve. *Autonomic Neuroscience*, 208, 66–72.
- Nobre, F., da Silva, C. A. A., Coelho, E. B., Salgado, H. C., & Fazan, R. (2006). Antihypertensive agents have different ability to modulate arterial pressure and heart rate variability in 2K1C rats. *American Journal of Hypertension*, 19, 1079–1083.
- Pereira-Junior, P. P., Chaves, E. A., Costa-e-Sousa, R. H., Masuda, M. O., Campos de Carvalho, A. C., & Nascimento, J. H. M. (2006). Cardiac autonomic dysfunction in rats chronically treated with anabolic steroid. *European Journal of Applied Physiology*, 96, 487–494.
- Pereira-Junior, P. P., Marocolo, M., Rodrigues, F. P., Medei, E., & Nascimento, J. H. M. (2010). Noninvasive method for electrocardiogram recording in conscious rats: Feasibility for heart rate variability analysis. *Anais da Academia Brasileira de Ciências*, 82, 431–443.
- Powell, K. L., Liu, Z., Curl, C. L., Raaijmakers, A. J. A., Sharma, P., Braine, E. L., Gomes, F. M., Sivathamboo, S., Macefield, V. G., Casillas-Espinosa, P. M., Jones, N. C., Delbridge, L. M., & O'Brien, T. J. (2021). Altered cardiac structure and function is related to seizure frequency in a rat model of chronic acquired temporal lobe epilepsy. *Neurobiology of Disease*, 159, 105505.
- Quagliotto, E., Casali, K. R., Dal Lago, P., & Rasia-Filho, A. A. (2015). Neuropeptides in the posterodorsal medial amygdala modulate central cardiovascular reflex responses in awake male rats. *Brazilian Journal of Medical and Biological Research*, 48, 128–139.
- Ramaekers, D., Beckers, F., Demeulemeester, H., & Aubert, A. E. (2002). Cardiovascular autonomic function in conscious rats: A novel approach to facilitate stationary conditions. *Annals of Noninvasive Electrocardiology*, 7, 307–318.
- Ribeiro, A. B., da Silva, T. M., Santos-Junior, N. N., Castania, J. A., Fayán, R., Jr., & Salgado, H. C. (2021). Short-term effect of ligature-induced periodontitis on cardiovascular variability and inflammatory response in spontaneously hypertensive rats. *BMC Oral Health*, 21, 515.
- Rowan, W. H., III, Campen, M. J., Wichers, C. L., & Watkinson, W. O. (2007). Heart rate variability in rodents: Uses and caveats in toxicological studies. *Cardiovascular Toxicology*, 7, 28–51.
- Ruiz, R., Ramos, S. P., Pinge, M. M., de Moraes, S. F., & Polito, M. (2014). Caffeine and physical training: Effects on cardiac morphology and cardiovascular response. *Revista da Associação Médica Brasileira*, 60, 23–28.
- Saalfeld, J., & Spear, L. (2014). Developmental differences in the effects of alcohol and stress on heart rate variability. *Physiology and Behavior*, 135, 72–80.
- Sallam, M. Y., El-Gowilly, S. M., Abdel-Galil, A. G. A., & El-Mas, M. M. (2016). Central GABA_A receptors are involved in inflammatory and cardiovascular consequences of endotoxemia in conscious rats. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 389, 279–288.
- Sallam, M. Y., El-Gowilly, S. M., Abdel-Galil, A. G. A., & El-Mas, M. M. (2017). Cyclosporine counteracts endotoxemia-evoked reductions in blood pressure and cardiac autonomic dysfunction via central sGC/MAPKs signaling in rats. *European Journal of Pharmacology*, 797, 143–152.
- Sant'Ana, J. E., Pereira, M. G. A. G., Da Silva, J. D., Dambrós, C., Costa-Neto, C. M., & Souza, C. D. (2011). Effect of the duration of daily aerobic physical training on cardiac autonomic adaptations. *Autonomic Neuroscience*, 159, 32–37.
- Sanyal, S. N., Arita, M., & Ono, K. (2002). Inhomogeneous derangement of cardiac autonomic nerve control in diabetic rats. *Circulation Journal*, 66, 283–288.
- Scridon, A., Gallet, C., Arisha, M. M., Oréa, V., Chapuis, B., Li, N., Tabib, A., Christé, G., Julien, C., & Chevalier, P. (2012). Unprovoked atrial tachyarrhythmias in aging spontaneously hypertensive rats: The role of the autonomic nervous system. *American Journal of Physiology. Heart and Circulatory Physiology*, 303(3), H386–H392.
- Shi, S., Liu, T., Wang, D., Zhang, Y., Liang, J., Yang, B., & Hu, D. (2017). Activation of N-methyl-D-aspartate receptors reduces heart rate variability and facilitates atrial fibrillation in rats. *EP Europace*, 19, 1237–1243.
- Shi, Z. J., Cheng, M., Liu, Y. C., Fan, X. R., & Wei, Z. Y. (2020). Effect of chronic intermittent hypobaric hypoxia on heart rate variability in conscious rats. *Clinical and Experimental Pharmacology and Physiology*, 47, 60–66.
- Silva, L. E. V., Geraldini, V. R., de Oliveira, B. P., Silva, C. A. A., Porta, A., & Fazan, R. (2017). Comparison between spectral analysis and symbolic dynamics for heart rate variability analysis in the rat. *Scientific Reports*, 7, 8428.
- Silva, L. E. V., Silva, C. A. A., Salgado, H. C., & Fazan, R., Jr. (2017). The role of sympathetic and vagal cardiac control on complexity of heart rate dynamics. *American Journal of Physiology-Heart and Circulatory Physiology*, 312, H469–H477.
- Simoes, M. R., Azevedo, B. F., Fiorim, J., Freire, D. D. J., Covre, E. P., Vassallo, D. V., & dos Santos, L. (2016). Chronic mercury exposure impairs the sympathovagal control of the rat heart. *Clinical and Experimental Pharmacology and Physiology*, 43, 1038–1045.
- Soler, A. I. R., Silva, L. E. V., Fayán, R., Jr., & Murta, L. O., Jr. (2018). The impact of artifact correction methods of RR series on heart rate variability parameters. *Journal of Applied Physiology*, 124, 646–652.
- Svorc, P., Jr., Svorc, P., & Gresova, S. (2023). Sex differences, chronobiology and general anesthesia in activities of the autonomic nervous system in rats. *Experimental Physiology*, 108, 1–8.
- Towa, S., Kuwahara, M., & Tsubone, H. (2004). Characteristics of autonomic nervous function in Zucker-fatty rats: Investigation by power spectral analysis of heart rate variability. *Experimental Animals*, 53, 137–144.
- Tsai, S. H., Lin, J. Y., Lin, Y. C., Liu, Y. P., & Tung, C. S. (2020). Portal vein innervation underlying the pressor effect of water ingestion with and without cold stress. *The Chinese Journal of Physiology*, 63, 53–59.

- Wang, D., Liu, T., Shi, S., Li, R., Shan, Y., Huang, Y., Hu, D., & Huang, C. (2016). Chronic administration of catestatin improves autonomic function and exerts cardioprotective effects in myocardial infarction rats. *Journal of Cardiovascular Pharmacology and Therapeutics*, 21, 526–535.
- Xu, L., Zhou, L., Dai, J., & Lin, Y. (2020). Effects of guanxinling tablets on the autonomic nerve regulation function of MI/RI rats by antioxidation and inhibition of inflammatory response. *Acta Medica Mediterranea*, 36, 3389–3394.
- Yang, Y. N., Tsai, H. L., Lin, Y. C., Liu, Y. P., & Tung, C. S. (2019). Differential effects of sympatholytic agents on the power spectrum of rats during the cooling-induced hemodynamic perturbations. *Chinese Journal of Physiology*, 62, 86–92.
- Zajackowski, S., Smolińska, M., Badtke, P., & Wierzbę, T. H. (2014). Time-domain and spectral analysis of heart rate variability in rats challenged with hypoxia. *Computers in Cardiology*, 41, 785–788.
- Zajackowski, S., Ziolkowski, W., Badtke, P., Zajackowski, M. A., Flis, D. J., Figarski, A., Smolińska-Bylańska, M., & Wierzbę, T. H. (2018). Promising effects of xanthine oxidase inhibition by allopurinol on autonomic heart regulation estimated by heart rate variability (HRV) analysis in rats exposed to hypoxia and hyperoxia. *PLoS ONE*, 13(2), e0192781.
- Zhu, G., Chen, Z., Dai, B., Zheng, C., Jiang, H., Xu, Y., Sheng, X., Guo, J., Dan, Y., Liang, S., & Li, G. (2018). Chronic lead exposure enhances the sympathoexcitatory response associated with P2X4 receptor in rat stellate ganglia. *Environmental Toxicology*, 33, 631–639.

How to cite this article: Švorc, P. Jr, Grešová, S., & Švorc, P. (2023). Heart rate variability in male rats. *Physiological Reports*, 11, e15827. <https://doi.org/10.14814/phy2.15827>